Grief from losing a loved one is a normal experience that, unfortunately, everyone goes through at some point. Grief is the most common experience humans have, but that reality doesn't help the emotional, psychological, and physical pain the death of a loved one can bring. Days stuck on a spinning wheel of thoughts: "What if I had just..?" "Why didn't I..?" "I wish I had..." And nights tormented by dreams, images and memories of the deceased, and sudden waking to the reality that they are really gone; the moments between sleep and wake can become the most dreaded time of the day.

Losing a loved one is hard enough when it's "natural" or "expected," such as due to old age or an extended battle with cancer. If the death was traumatic in some way, such as sudden, violent, suicide, homicide, or if the person was young, there are a multitude of additional stressors that can compound the psychological experience.

Here are some common experiences when it comes to grief:

- At first, feeling like it can't be real.
- Overwhelming waves of sadness, loss, longing, and crying that seem to come on at random times and leave just as suddenly. You may think it's done with, but then it returns and each time, you have to confront the loss all over again. These waves of grief tend to be at their highest intensity for several months following the death, and tend to lessen on their own with time, with spikes along the way.
- · Seeking comfort in various ways.
- B yo th ofa a

Grieving the eath of a Loved One

Here are some ways to help yo irself:

- Don't make major decisions or life changes when you're in the sharp pangs of grief.
- Give it time. This one's a cliche for a reasun. Time I eally does help ease the physical and emotional waives of grief. We are naturally wired toward healing.
- Cry it out. It's natural for humans to feel the urge to cry when sad, so when it comes on, let the tears flow. (Science fad: Crying allows the body to literally release stress hormones through tears.) Ele ause the sudden waves of grief can show up at any point in the day or night, at metimes people isolate in anticipation of being embarrassed if the uncontrollable urge to try comes on in public. There is no need to isolate or explain yourself (although sometimes disclosing to someone near you what's happening can actually force surprising new friendships). Another valid diche: The ones who matter don't mind; the cines who mind don't matter. You might want to bring hissues when you go out in public, just in case, and locate the nearest bathroom if you start to feel the urge to cry and prefer to seek temporary private solace.
- View and treat yourself a ; you would you it est frie id in the same situation. What would you tell them if they were experien ing the same circumstances as you are now?
- Normalize your thoughts and emotions. The relisting first way to grieve, and what you experience during grief varies depend ing on the specific dircumstances of the death, your own life circumstances, and you rivelation to the deceased, among other factors. Resist judging your emotions as good or had when you feel them; instead, practice observing their ebb and it woulth kindness and compassion for yourself. Remind yourself that you are no alone in how you feel. The strong likelihood is if you're having a thought or leaking, in all of human history others have had/are having it, to Podcasts, blog, and gief support groups are great ways to be reminded of this, which can be a sally or inforting. You are not alone in how you're thinking and seling...ever.
- Keep on a regular routine of eating/nutrit on, hygiene, exercise, hydration, and sleep, even if it doesn't siem enticing or in a portant in the moment. The following are general guidelines for taking care of y x rself, it at a ways check with your doctor and follow medica advice specific to you. Eat 3: quare meals a day plus healthy snacks between. Exercise 30 min It is a day in a way that works for you; even long walks can be in credibly helpful f you don't fe al like doing your usual cardio or strength trainin j. Go to bed earl / and alle t ye arself the full 8 hours, even if your sleep quality suck ; for a while. Do not these him is that you know are good for you, even-- especially -- when you dor t feel like it, nay be difficult when you're grieving. But once you're doing them, I thir k you'll be unprised by how much of a difference it makes in ho vyou feel. The r x re irrat ona thoughts that are often part of grief-- for example, blaming yours all for a level one's death when it really had nothing to do with you -- can get stroke you're tired, unkempt, hungry, or dehydrated. You may tart to notice this pattern for yourself. For example, you know that when your mind starts to go hay vire with in ational or morbid thoughts and it's 9:00 PIVI it may e time for bed.
- Use healthy coping skills that give you con ort and joy or something approximating the joy you knew before you to loss-- and remind you that not all is bad in the world and the plane still things to look forward door. What were our general coping mechanism his prior to the loss of a loved one become our amplified go-tos in times of distres in some way. These might include deep breathing to reditating, positive affirmations (see the end of this article for a list related to to perform any to making music, journaling, writing poetry/stories, or obtaining, making a to listening to making music, journaling, interacting with friends and family, being in nature and taking care of a pet, among others.



- Develop thought boundaries with yourself so that grief doesn't take over your whole life. You are still alive, and you deserve to live a good life, as painful as that may be to acknowledge at times when you're grieving. This might look like scheduling a certain amount of daily or weekly time to talk and think about your loved one, then moving on with your day knowing you have already planned more time to devote to them. Being in therapy can provide a great way to practice thought boundaries, because you can decide with your therapist that you will, for example, practice focusing on your deceased loved one only during therapy sessions and related homework exercises. Or, it could be simply determining where your limit is in terms of your thoughts; for example, maybe you decide in advance that once you're repeating the same distressing thought over and over without resolution (ruminating), you know that it's time to take action to resolve the thought or to let it go and do something positive.
- If you had conflict with the deceased, or if they hurt you, work toward forgiving them. If the hurt was so bad that you don't think you can or should forgive them (people do heinous things sometimes), focus instead on the benefits of forgiveness for you, not them. Therapy can help you work through this process.
- Keep it specific to this experience. Sometimes a sad or traumatic event can cause
 us to generalize our experiences into global beliefs about people, life, and the
 world. In reality, both good and bad things happen in life, bad things sometimes
 happen to people who overwhelmingly do good with their lives, and good things
 sometimes happen to people who overwhelmingly do bad. Life can really suck and
 be unfair sometimes, and it can also be mundane or wonderful at other times.

When should you seek professional help?

• First and foremost, as mentioned above, you can seek therapy at any point in time. It is not just for obvious "problems," but also intended for processing experiences



